## Osmolite 1.5kcal 1.5 kcal/ml complete, balanced liquid

#### FOR HEALTHCARE PROFESSIONAL USE ONLY

#### **PRESENTATION**

- Presented in 500 ml (754 kcal), 1000 ml (1507 kcal) & 1500 ml (2261 kcal) Ready to Hang (RTH) containers.
- · Osmolite 1.5kcal is unflavoured.

#### **USES**

Food for Special Medical Purposes, for use under medical supervision. Suitable as a sole source of nutrition or as a nutritional supplement for patients who cannot or will not eat sufficient quantities of everyday food and drink to meet their nutritional requirements.

Nutritionally complete for vitamins and minerals in 1071 ml (excluding electrolytes, calculated using the UK Reference Nutrient Intake for men aged 19-50 years).

#### COMMUNITY USE—PRESCRIPTIONS

Can be prescribed on a FP10 (GP10 in Scotland) for the following indications:

- · Disease-related malnutrition
- Total gastrectomy
- Short bowel syndrome
- Bowel fistulae
- Intractable malabsorption
- Pre-operative preparation of patients who are malnourished
- Proven inflammatory bowel disease
- Dvsphagia
- CAPD
- Haemodialysis

All prescriptions should be endorsed ACBS (Advisory Committee on Borderline Substances).

Available on the DPS (Drugs Payment Scheme) and GMS (General Medical Services) scheme in Ireland.

#### **STORAGE**

- Store unopened at room temperature.
- Avoid prolonged exposure to light.
- Once opened, unused product should be resealed and stored in a refrigerator.
- Unused contents should be discarded after 24 hours.

#### DIRECTIONS FOR USE

- Ready for use.
- Administer at room temperature for tube feeding.
- The volume/flow rate should be adjusted to meet the patient's nutritional needs and tolerance. This product has a low viscosity and will pass down a fine nasogastric tube.
- A Flexitainer enteral nutrition container may be used if decanting is necessary.
- An Abbott enteral feeding pump may be used in conjunction with the Abbott enteral feeding system where a more accurately controlled delivery of feed is indicated. An ambulatory system is available for use.
- For gravity feeding, the use of a Flexiflo gravity gavage set is recommended.

#### **PRECAUTIONS**

- In patients receiving some medications there may be a risk of drug nutrient interactions (e.g. warfarin and vitamin K). Careful prescribing and monitoring practices will serve to reduce the risk (please refer to pharmacist).
- Patients should not make any additions to the feed without consulting their pharmacist or dietitian.
- Unless recommended by a qualified healthcare professional, not intended for use in children.
- When feeding to patients with dysphagia, please thicken the product as appropriate.

#### **CONTRA-INDICATIONS**

- FOR ENTERAL USE ONLY.
- Do not use in children under 1 year of age.
- Not for use in galactosaemia.
- Suitable for people with diabetes provided that routine glucose checks are performed.

#### **INGREDIENTS**

Water, maltodextrin, *milk* proteins, vegetable oils (high oleic sunflower, canola, MCT from palm kernel oil), *soy* protein isolate, minerals (potassium citrate, calcium phosphate, tribasic, magnesium chloride, sodium citrate, potassium chloride, potassium phosphate dibasic, ferrous sulphate, zinc sulphate, manganese sulphate, cupric sulphate, sodium molybdate, chromium chloride, sodium selenate, potassium iodide), stabilisers (E460, E466, E418), emulsifier: *soy* lecithin, choline chloride, vitamins (C, E, niacinamide, calcium pantothenate, B<sub>6</sub>, B<sub>1</sub>, B<sub>2</sub>, vitamin A palmitate, beta carotene, folic acid, K<sub>1</sub>, biotin, D<sub>3</sub>, B<sub>12</sub>), taurine, l-carnitine.

#### GENERAL INFORMATION

Energy density	1.5 kcal/ml	
Energy distribution: Protein Carbohydrate Fat	16.6% 54.0% 29.3%	
Renal solute load	501 mOsm/L	
Osmolarity	392 mOsm/L	
Osmolality	$510 \text{ mOsm/kg H}_2\text{O}$	
Gluten free?	✓	
Clinically lactose free?	✓	
Milk free?	×	
Suitable for vegetarians?	<b>√</b> 1	

For suitability for other diets and free-from information, please contact the Freephone Nutrition Helpline on 0800 252882.

 Vitamin D is synthesised from cholesterol, extracted from the grease in wool sheared from live sheep.



# Osmolite 1.5kcal 1.5 kcal/ml complete, balanced liquid

### FOR HEALTHCARE PROFESSIONAL USE ONLY

NUTRITION INFORMATION			
	units	per 100 ml	_
Energy	kJ	634	
	kcal	151	
Fat	g	4.91	
- of which saturates	g	1.3	
- of which MCT*	g	0.9	
Carbohydrate	g	20.4	
- of which sugars	g	1.15	
Protein (nitrogen)	g	6.27 (1.0)	
Salt	g	0.35	
Fibre	g	О	
***			
Vitamins		160	
Vitamin A (RE) - of which β-carotene	μg	160	
Vitamin D <sub>3</sub>	μg	34.5 1.00	
Vitamin E ( $\alpha$ TE)	μg mg	2.6	
Vitamin K <sub>1</sub>	μg	8.0	
Vitamin C		13	
Folacin (folic acid)	mg μg	40	
Thiamin (vitamin B <sub>1</sub> )	mg	0.26	
Riboflavin (vitamin B <sub>2</sub> )	mg	0.34	
Vitamin B <sub>6</sub>	mg	0.39	
Vitamin B <sub>12</sub>	μg	0.40	
Niacin (NE)	mg	2.9	
Pantothenic acid	mg	1.3	
Biotin	μg	7.2	
Minerals			
Sodium	mg (mmol)	140 (6.09)	
Potassium	mg (mmol)	165 (4.22)	
Chloride	mg (mmol)	145 (4.09)	
Calcium	mg (mmol)	100 (2.50)	
Phosphorus (phosphate)	mg (mmol)	100 (3.23)	
Magnesium	mg (mmol)	31 (1.28)	
Iron	mg	2.2	
Zinc	mg	1.9	
Manganese	mg	0.50	
Copper	mg	0.25	
Iodine	μg	15	
Selenium	μg	7.0	
Chromium	μg	10.0	
Molybdenum		15	
Choline	μg	60	
L-carnitine	mg mg	12	
Taurine	mg mg	15	
Taurine	mg	±0	
Water	g	77.2	
	0	, , . <del> _</del>	

PROTEIN & AMINO ACIDS			
	g/100 g protein	g/100 ml	
Protein source			
Sodium caseinate	73.0	4.58	
Soy protein isolate	16.0	1.00	
Calcium caseinate	11.0	0.69	
Amino acids			
- Essential			
Histidine	2.94	0.18	
Isoleucine	5.45	0.34	
Leucine	9.50	0.60	
Lysine	7.85	0.49	
Methionine	2.73	0.17	
Phenylalanine	5.28	0.33	
Threonine	4.33	0.27	
Tryptophan	1.15	0.07	
Valine	6.72	0.42	
Arginine	4.41	0.28	
-Non-essential			
Alanine	3.25	0.20	
Aspartic acid	3.32	0.21	
Cystine	0.67	0.04	
Glutamic acid	11.7	0.73	
Glycine	2.27	0.14	
Proline	9.85	0.62	
Serine	5.91	0.37	
Tyrosine	5.48	0.34	
Asparagine	4.58	0.29	
Glutamine	10.1	0.63	
Non-protein calorie: N	125 :1		

CARBOHYDRATES			
	% total carbohydrates	g/100 ml	
Carbohydrate source			
Maltodextrin	99.3	20.3	
Carboxymethyl cellulose	0.66	0.14	

FAT & FATTY ACIDS			
		% total fatty acids	g/100 ml
Fat source			
High alais and arran a	:1	40.4	2.00
High oleic sunflower of	11	48.4	2.38
	1 - 9	29.0	1.42
MCT from palm kernel Lecithin	1 011	19.3	0.95
Lecitiiiii		3.30	0.16
Fatty acids		g/100 g fat	g/100 ml
- Essential		0/ 0	0/
Linoleic acid	C18:2	11.1	0.55
Alpha-linolenic acid	C18:3	2.67	0.13
· · · · · · · · · · · · · · · · · · ·		, ,	<u> </u>
- Monounsaturated			
Palmitoleic acid	C16:1	0.09	trace
Oleic acid	C18:1	55.5	2.77
Gadoleic acid	C20:1	0.50	0.02
Erucic acid	C22:1	0.21	0.01
- Saturated			
Caproic acid	C6:0	0.09	trace
Caprylic acid	C8:0	10.5	0.52
Capric acid	C10:0	7.39	0.37
Lauric acid	C12:0	0.23	0.01
Myristic acid	C14:0	0.09	trace
Palmitic acid	C16:0	3.54	0.18
Margaric acid	C17:0	-	-
Stearic acid	C18:0	2.09	0.10
Arachidic acid	C20:0	0.34	0.02
Behenic acid	C22:0	0.51	0.03
Tricosanoic acid	C23:0	-	-
Lignoceric acid	C24:0	0.17	0.01
_	-		
P/S ratio	0.56		

Abbott Laboratories Ltd., Abbott House, Vanwall Business Park, Vanwall Rd., Maidenhead, Berks, SL6 4XE. Tel: 0800 252882 Nutrition.abbott/uk

4.9: 1

Abbott Laboratories (Ireland) Ltd., Liffey Valley Office Campus, Dublin 22. Tel: (01) 4691500 Fax: (01) 4691501 abbottnutrition.ie

n6:n3 ratio

ANUKANI180312 Date of preparation: November 2018 Version 4

