## **PRESENTATION**

- Presented in 500 ml (759 kcal) Ready to Hang (RTH) bottles.
- Available in vanilla flavour. NB There is a separate datasheet for PaediaSure Plus Fibre sip feed.

## **USES**

Food for Special Medical Purposes, for use under medical supervision. Suitable as a nutritional supplement or as a sole source of nutrition for patients who cannot or will not eat sufficient quantities of everyday food and drink to meet their nutritional requirements.

Nutritionally complete for vitamins and minerals in 619 ml for children aged 1-3 years, 667 ml for children aged 4-6 years and 833 ml for children aged 7-10 years (excluding electrolytes, calculated using the UK Reference Nutrient Intake for these age bands).

#### COMMUNITY USE—PRESCRIPTIONS

Can be prescribed on a FP10 (GP10 in Scotland) for the following indications in children weighing 8-30 kg:

- Disease-related malnutrition and/or growth failure
- Short bowel syndrome
- Bowel fistulae
- Intractable malabsorption
- Pre-operative preparation of patients who are malnourished
- Dysphagia

All prescriptions should be endorsed ACBS (Advisory Committee on Borderline Substances).

#### STORAGE

- Store unopened at room temperature.
- Avoid prolonged exposure to light.
- Once opened, unused product should be resealed and stored in a refrigerator.
- Unused contents should be discarded after 24 hours.

# DIRECTIONS FOR USE

- Ready for use.
- Administer at room temperature for tube feeding.
- The volume/flow rate should be adjusted to meet the patient's nutritional needs and tolerance. This product has a low viscosity and will pass down a fine nasogastric tube.
- An Abbott enteral feeding pump may be used in conjunction with the Abbott enteral feeding system where a more accurately controlled delivery of feed is indicated. An ambulatory system is available for use.
- For gravity feeding, the use of a Flexiflo gravity gavage set is recommended.
- A Flexitainer enteral nutrition container may be used if decanting is necessary.

#### **PRECAUTIONS**

- In patients receiving some medications there may be a risk of drug nutrient interactions (e.g. warfarin and vitamin K). Careful prescribing and monitoring practices will serve to reduce the risk (please refer to pharmacist).
- Patients should not make any additions to the feed without consulting their pharmacist or dietitian.
- Many nutritional products contain sucrose and other sugars. It is important for patients who are taking supplements as sip feeds to observe good oral hygiene. It is suggested that patients consult with their dentist for further advice.
- When feeding to patients with dysphagia, please thicken the product as appropriate.

## CONTRA-INDICATIONS

- FOR ENTERAL USE ONLY.
- Do not use in children under 1 year of age.
- Not for use in galactosaemia.
- Suitable for people with diabetes provided that routine glucose checks are performed.

# **INGREDIENTS**

Water, maltodextrin, vegetable oils (high oleic sunflower, soy, MCT from palm kernel oil), *milk* proteins, sucrose, minerals (potassium citrate, magnesium chloride, calcium phosphate tribasic, potassium phosphate dibasic, potassium chloride, ferrous sulphate, zinc sulphate, cupric sulphate, manganese sulphate, sodium fluoride, potassium iodide, sodium molybdate, chromium chloride, sodium selenite), *oat* fibre, fructooligosaccharides, *soy* polysaccharide, flavouring, emulsifiers (E471, E322: *soy* lecithin), gum arabic, carboxy methylcellulose, choline bitartrate, vitamins (C, E, niacinamide, calcium pantothenate, vitamin A palmitate, B<sub>2</sub>, B<sub>1</sub>, B<sub>6</sub>, D<sub>3</sub>, folic acid, biotin, K<sub>1</sub>, B<sub>12</sub>), myo-inositol, taurine, l-carnitine.

#### GENERAL INFORMATION

Energy density	1.5 kcal/ml
Energy distribution: Protein Carbohydrate Fat Fibre (FOS)	11.1% 43.2% 44.3% 1.45%
Renal solute load	354 mOsm/L
Osmolarity	270 mOsm/L
Osmolality	347 mOsm/kg H₂O
Gluten free?	✓
Clinically lactose free?	✓
Milk free?	×
Suitable for vegetarians?	<b>√</b> 1
Suitable for Halal diet?	✓
Suitable for Kosher diet?	✓

For further free-from information, please contact the Freephone Nutrition Helpline on 0800 252882.

Vitamin D is synthesised from cholesterol, extracted from the grease in wool sheared from live sheep.

Version 6: June 2019



\*fructo-oligosaccharides

# 1.5 kcal/ml complete, balanced nutrition with fibre and FOS\* for children weighing 8-30 kg

#### **NUTRITION INFORMATION** units per 100 ml Energy 635 kcal 152 7.47 - of which saturates g 2.0 - of which MCT\*\* 1.37 Carbohydrate 16.4 g - of which sugars 2.2 Fibre g - of which FOS\* 0.35 Protein (nitrogen) g 4.20 (0.67) Salt 0.15 Vitamins Vitamin A (RE) μg 99 - of which β-carotene μg O Vitamin D<sub>3</sub> μg 1.13 Vitamin E (α TE) mg 1.5 Vitamin K<sub>1</sub> μg 5.9 Vitamin C mg 7.5 Folacin (folic acid) μg 23 Thiamin (vitamin B<sub>1</sub>) mg 0.23 Riboflavin (vitamin B2) mg 0.30 Vitamin B<sub>6</sub> mg 0.15 Vitamin B<sub>12</sub> 0.30 μg Niacin (NE) mg 1.8 Pantothenic acid mg 0.45 Biotin μg 7.5 Minerals Sodium mg (mmol) 60 (2.61) Potassium mg (mmol) 135 (3.45) Chloride mg (mmol) 100 (2.82) 83 (2.07) Calcium mg (mmol) Phosphorus (phosphate) mg (mmol) 80 (2.58) Magnesium mg (mmol) 24 (0.99)Iron mg 1.5 Zinc mg 1.5 Manganese 0.15 Copper mg 0.17 Iodine μg 15 Selenium μg 4.2 Chromium 3.8 μg Molvbdenum μg 5.9 Fluoride mg 0.08 Taurine 10.8 mg L-carnitine 2.6 mg Inositol 12 mg Choline mg 22.5

g

78.1

PKULEI	N & AMINO		
	g/100 g protein	g/100 ml	
Protein source			
Sodium caseinate	60.0	2.52	
Calcium caseinate	22.0	0.92	
Whey protein concentrate	18.0	0.76	
Amino acids			
- Essential			
Histidine	2.39	0.10	
Isoleucine	4.83	0.20	
Leucine	9.13	0.38	
Lysine	7.54	0.32	
Methionine	2.48	0.10	
Phenylalanine	4.54	0.19	
Threonine	4.80	0.20	
Tryptophan	1.21	0.05	
Valine	5.81	0.24	
Arginine	3.26	0.14	
- Non-essential			
Alanine	3.23	0.14	
Aspartic acid	3.18	0.13	
Cystine	0.73	0.03	
Glutamic acid	11.1	0.47	
Glycine	1.87	0.08	
Proline	9.69	0.41	
Serine	5.38	0.23	
Tyrosine	4.69	0.20	
Asparagine	4.45	0.19	
Glutamine	9.78	0.41	
Non-protein calorie: N	205:1		
Tion protein calone, it	200.1		

CARBOHYDRATES			
	% total carbohydrates	g/100 ml	
Carbohydrate source			
Maltodextrin	89.7	14.7	
Sucrose	10.0	1.64	
Oat fibre	0.11	0.02	
Fructo-oligosaccharide powder	0.09	0.01	
Soy fibre	0.05	trace	
Gum arabic	0.03	trace	
Carboxymethyl cellulose	0.01	trace	

FIBRE			
	% total fibre	g/100 ml	
Fibre source			
FOS* powder	31.1	0.34	
Oat fibre	31.0	0.34	
Soy fibre	20.7	0.23	
Gum arabic	12.1	0.13	
Carboxymethyl cellulose	5.16	0.06	
Soluble fibre content: 49%	Insoluble fibre content: 51%		

EAT & EATTV ACIDS

TIDDE

FAT & FATTY ACIDS				
		% total fatty acids	g/100 ml	
Fat source				
High oleic sunflower oi	1	49.5	3.70	
Soy oil		29.7	2.22	
MCT from palm kernel	oil	19.8	1.48	
Lecithin		1.07	0.08	
		,		
Fatty acids		g/100 g fat	g/100 ml	
- Essential				
Linoleic acid	C18:2	19.4	1.46	
Linolenic acid	C18:3	1.51	0.11	
- Monounsaturated				
Palmitoleic acid	C16:1	0.12	0.01	
Oleic acid	C18:1	46.3	3.49	
Petroselinic acid	C18:1	0.08	0.01	
Gadoleic acid	C20:1	0.05	trace	
Erucic acid	C22:1	-	-	
- Saturated				
Caproic acid	C6:0	0.09	0.01	
Caprylic acid	C8:o	10.5	0.79	
Capric acid	C10:0	7.38	0.56	
Lauric acid	C12:0	0.14	0.01	
Myristic acid	C14:0	0.06	trace	
Palmitic acid	C16:0	5.35	0.40	
Margaric acid	C17:0	0.07	0.01	
Stearic acid	C18:0	2.77	0.21	
Arachidic acid	C20:0	0.34	0.03	
Behenic acid	C22:0	0.56	0.04	
Tricosanoic acid	C23:0	-	-	
Lignoceric acid	C24:0	0.10	0.01	
P/S ratio	0.76			
n6:n3	12.8 : 1			

Abbott Laboratories Ltd., Abbott House, Vanwall Business Park, Vanwall Rd., Maidenhead, Berks. SL6 4XE. Tel: 0800 252882 Nutrition.abbott/uk

Abbott Laboratories (Ireland) Ltd., Liffey Valley Office Campus, Dublin 22. Tel: (01) 4691500 Fax: (01) 4691501 abbottnutrition.ie



Water

<sup>\*</sup> Fructooligosaccharides \*\*medium-chain triglycerides (C6:0 - C12:0)